

LSV Wintertriticale Ökolandbau 2012 - 2017, Anbaugebiet Sandstandorte Nordost

Kornertrag (86 % TS) dt/ha

	2017			
	Plöwen	Gülzow	Schmerwitz	Futterkamp
Land	MV	MV	BB	SH
Tulus	16,7	39,8	31,2	55,9
Securo	21,1	45,0	26,6	45,1
Rhenio	21,7	43,5	30,0	53,7
Salto		40,3	27,0	
Lombardo	19,9	44,9	29,2	44,8
Cedrico	17,9	48,6	28,9	45,9
Trefl	21,5	50,5	29,8	
N= 3	19,8	42,8	29,3	51,5
Grenzdifferenz (dt/ha)	2,5	8,7	1,7	4,9

2012-2017	
Sandstandorte Nordost	Anzahl Versuche
37,8	21
37,8	13
38,0	10
[]	5
[]	7
[]	4
[]	3
37,9	

Kornertrag (86 % TS) dt/ha relativ

	2017			
	Plöwen	Gülzow	Schmerwitz	Futterkamp
Land	MV	MV	BB	SH
Tulus	84	93	106	108
Securo	106	105	91	88
Rhenio	109	102	103	104
Salto		94	92	
Lombardo	100	105	100	87
Cedrico	90	114	99	89
Trefl	109	118	102	
100%=dt/ha	19,8	42,8	29,3	51,5
Grenzdifferenz rel.)	12,8	19,4	5,9	10,0
Grenzdifferenz (dt/ha)	2,5	8,7	1,7	4,9

2012-2017	
Sandstandorte Nordost	Anzahl Versuche
100	21
100	13
100	10
[]	5
[]	7
[]	4
[]	3
37,9	

Bezugsbasis = 100 % = Mittel Tulus, Securo, Rhenio